

## **POSSIBILITY OF UTILIZING SILKWORMS FOR MULTIPLICATION OF DENDROLIMUS SPECTABILIS CYTOPOLYHEDROSISVIRUS**

**R. Murakami and K. Miyamoto**

National Institute of Agrobiological Sciences, 1-2, Ohwashi, Tsukuba, Ibaraki 305-8634, Japan

E-mail: mritsuko@affrc.go.jp

### **ABSTRACT**

*Dendrolimus spectabilis* is the major pest of Pinus plants in East Asia, and *D. spectabilis* cytopolyhedrosis virus (DsCPV) has been used to control the pest in China and Japan. Large scale multiplication of DsCPV on its primary host *i.e.*, *D. spectabilis* works out expensive. Possibility of utilizing silkworms as an alternative host for multiplying DsCPV was explored as a cost effective measure. Thirty five races of silkworms were per orally inoculated with  $1 \times 10^8$  DsCPV polyhedral / g artificial diet. On examination after ten days, 29 races revealed the presence of DsCPV with the highest incidence of 26.7 % in Oh 1 Gou race. Hence, progenies of susceptible populations of Oh 1 Gou race were raised by subjecting to DsCPV inoculation followed by line breeding and directional selection for higher susceptibility for three generations. Resultant lines revealed a maximum DsCPV infection level of 90.0%. DsCPV obtained from the silkworms were confirmed by reverse transcription polymerase chain reaction method, and the presence of BmCPV if any, could also be differentiated.

**Key words:** *Dendrolimus spectabilis* cytopolyhedrosisvirus, DsCPV, infection, silkworm.